

# **STA Receives TGA Approval for ABRAXANE® in Combination with Gemcitabine for First-Line Treatment of Metastatic Pancreatic Cancer**

**Melbourne, Australia - March, 2014:** Australian biopharmaceutical company Specialised Therapeutics Australia (STA) is pleased to announce that ABRAXANE® (nanoparticle albumin-bound paclitaxel) in combination with gemcitabine is now approved by the Therapeutic Goods Administration (TGA) for the first-line treatment of metastatic pancreatic cancer.

The TGA approved indication is:

ABRAXANE, in combination with gemcitabine, is indicated for the first-line treatment of patients with metastatic adenocarcinoma of the pancreas.<sup>1</sup>

TGA approval was based on the pivotal randomised phase III trial, MPACT (Metastatic Pancreatic Adenocarcinoma Clinical Trial), published in the New England Journal of Medicine (NEJM) in October 2013.<sup>2</sup> The study reported that patients treated with ABRAXANE plus gemcitabine had a statistically significant improvement in overall survival (OS) compared to patients receiving the current standard of care, gemcitabine monotherapy (OS; median 8.5 months vs. 6.7 months; HR 0.72, P<0.001).<sup>2</sup> An updated analysis of OS presented at a recent international cancer conference in January 2014 (ASCO GI) showed the survival benefit was further extended in the ABRAXANE plus gemcitabine arm, with a 2.1 month median OS improvement compared to gemcitabine alone (OS; median 8.7 months vs 6.6 months; HR=0.72; p<0.0001).<sup>3</sup>

MPACT is the first phase III trial in metastatic pancreatic cancer to report greater than 3-year survival rates, with 4% of patients in the ABRAXANE plus gemcitabine arm alive after three years, and 3% of patients alive at 42 months,

compared to 0% in the gemcitabine alone arm at both time points.<sup>3</sup>

STA Chief Executive Officer Mr Carlo Montagner said TGA approval paves the way for Australian patients with metastatic pancreatic cancer to access this more effective treatment option.

He commented: “In Australia, pancreatic cancer is the fifth most common cause of death from cancer for both men and women, and very few treatment options exist for this group of patients. No new drugs have been approved by the TGA for this disease since 2006. We are extremely pleased to receive TGA approval in recognition that ABRAXANE is capable of prolonging survival for patients with metastatic pancreatic cancer, and look forward to a Pharmaceutical Benefits Scheme (PBS) listing for this difficult to treat cancer.”

ABRAXANE is now TGA approved for three indications; metastatic breast cancer, first-line Non-Small Cell Lung Cancer (NSCLC) and first-line metastatic pancreatic cancer.<sup>1</sup>

STA is currently seeking a PBS listing for ABRAXANE in first line metastatic pancreatic cancer.

## **About MPACT<sup>2</sup>**

MPACT (Metastatic Pancreatic Adenocarcinoma Clinical Trial), was a Celgene-sponsored, open-label, randomised, international study of 861 patients with metastatic pancreatic cancer. Patients were randomised to receive either ABRAXANE plus gemcitabine (125 mg/m<sup>2</sup> followed by 1000 mg/m<sup>2</sup> gemcitabine for 3 weeks followed by a week of rest) or gemcitabine alone (1000 mg/m<sup>2</sup> administered weekly for 7 weeks followed by a week of rest then weekly administration for 3 weeks followed by one week of rest).

The primary endpoint of the study was overall survival. Secondary endpoints were progression-free survival and overall response rate determined by independent radiological review. Other endpoints included the safety and tolerability of this combination in patients with metastatic pancreatic cancer.

The most common grade  $\geq 3$  treatment-related adverse events in MPACT for ABRAXANE plus gemcitabine vs. gemcitabine alone were neutropenia (38% vs. 27%), fatigue (17% vs. 7%), and peripheral neuropathy (17% vs. 1%) respectively. The median time to neuropathy improvement by one grade from grade  $\geq 3$  was 21 days in the ABRAXANE plus gemcitabine arm compared to 29 days in the gemcitabine alone arm. Neuropathy improved to grade 1 or lower in a median of 29 days for the ABRAXANE plus gemcitabine arm and was not reached for the gemcitabine alone arm. There was no difference in serious life threatening toxicity (4% in each arm).<sup>2</sup>

## About Advanced Pancreatic Cancer<sup>4</sup>

Advanced pancreatic cancer is a difficult-to-treat cancer with the lowest survival rates among all cancer types. Across all patients with pancreatic cancer, relative 5-year survival is approximately 5% and is less than 2% for those with advanced disease. There are two main types of pancreatic cancer - adenocarcinomas, which accounts for approximately 90% of all pancreatic cancer, and neuroendocrine tumours. Pancreatic cancer is relatively uncommon with new cases accounting for only 2.1% of all newly diagnosed cancers. However, pancreatic cancer is the fifth most common cause of cancer death for men and women in the United States and Australia, and the ninth most commonly diagnosed cancer in Australia.<sup>4</sup>

## About ABRAXANE<sup>®</sup>

Developed using the proprietary *nab*<sup>™</sup> technology platform, ABRAXANE is a nanoparticle protein-bound chemotherapy agent. ABRAXANE combines paclitaxel with albumin, a naturally-occurring human protein, to deliver the drug and therefore eliminates the need for solvents in the administration process. Nanoparticle technology allows ABRAXANE to deliver a 49% higher dose compared to regular solvent-based paclitaxel without compromising safety and tolerability.<sup>1,5</sup>

ABRAXANE is approved for the treatment of metastatic breast cancer, NSCLC and metastatic pancreatic cancer.<sup>1</sup> In Australia, ABRAXANE is currently listed on the PBS for the treatment of metastatic breast cancer and HER2 positive breast cancer in combination with trastuzumab. Abraxane is not PBS listed for the indications of NSCLC or metastatic pancreatic cancer.

For the first-line treatment of metastatic pancreatic cancer, the recommended dosing regimen for ABRAXANE is 125 mg/m<sup>2</sup> administered as an intravenous infusion over 30 minutes on Days 1, 8 and 15 of each 28-day cycle. Gemcitabine 1000 mg/m<sup>2</sup> is administered as an intravenous infusion beginning immediately after the completion of ABRAXANE administration on Days 1, 8 and 15 of each 28-day cycle.<sup>1</sup>

ABRAXANE is currently in various stages of investigation for the treatment of the following cancers: metastatic melanoma, adjuvant pancreatic cancer, bladder cancer and expanded applications for breast cancer.

BEFORE PRESCRIBING PLEASE CONSULT THE ABRAXANE PRODUCT INFORMATION AVAILABLE AT [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

## **ABRAXANE®          Minimum          Product** **Information**

ABRAXANE: Nanoparticle albumin-bound paclitaxel 100 mg powder for injection (suspension)

### **Indications:**

Metastatic carcinoma of the breast after failure of anthracycline therapy.

First-line treatment of non-small cell lung cancer (NSCLC) in combination with carboplatin, in patients who are not candidates for potentially curative surgery and/or radiation.

First-line treatment of patients with metastatic adenocarcinoma of the pancreas, in combination with gemcitabine.

**Contraindications:** Baseline neutrophil count  $< 1.5 \times 10^9/L$ , hypersensitivity to ABRAXANE or albumin, pregnancy, lactation.

**Precautions:** Administer under the supervision of a physician experienced in the use of chemotherapeutic agents. ABRAXANE is not clinically interchangeable with other paclitaxel formulations. Dose dependent and dose limiting bone marrow suppression (frequent peripheral blood cell counts recommended for all patients). Peripheral neuropathy, sepsis, severe hypersensitivity, pneumonitis, patients with hepatic impairment, cardiotoxicity, affects fertility, pregnancy (category D), lactation, paediatric use. In elderly - more frequent myelosuppression, peripheral neuropathy, arthralgia, diarrhoea, decreased appetite, dehydration and epistaxis. Refer to full PI for more information.

**Interactions:** Inhibitors or inducers of either CYP2C8 or CYP3A4 (e.g. inhibitors: erythromycin, ketoconazole, fluoxetine, imidazole antifungals, gemfibrozil, cimetidine, ritonavir, saquinavir, indinavir, and nelfinavir; inducers: rifampicin, carbamazepine, phenytoin, efavirenz, nevirapine). CYP2C8 and CYP3A4 substrates, quinidine, PEG-35 castor oil, quercetin, clozapine, morin, and resveratrol. Refer to full PI for details.

**Adverse Effects:** Very common effects in ABRAXANE monotherapy: Neutropenia, anaemia, leukopenia, thrombocytopenia, lymphopenia, bone marrow suppression, peripheral neuropathy, neuropathy, hypoaesthesia, paraesthesia, myalgia, arthralgia, asthenia, nausea, vomiting, diarrhoea, constipation, stomatitis, anorexia, pyrexia, alopecia, rash, fatigue, mucositis. Additional very common effects in combination with carboplatin: Peripheral oedema, dyspnoea, decreased appetite. Additional very common effects in combination with gemcitabine: chills, abdominal pain, dysgeusia, headache, dizziness, dehydration, hypokalemia, cough, epistaxis, weight decreased, ALA increased, pain in extremity, insomnia, depression, anxiety. This is not a full list of adverse effects - refer to full PI for more information.

**Dose:**

Metastatic Breast Cancer: ABRAXANE  $260 \text{ mg/m}^2$  every 3 weeks.

NSCLC: ABRAXANE  $100 \text{ mg/m}^2$  on days 1, 8, and 15 of each 21-day cycle. Recommended carboplatin dose is  $\text{AUC} = 6 \text{ mg} \cdot \text{min/mL}$  on day 1 only of each 21-

day cycle, beginning immediately after the end of ABRAXANE administration.

Metastatic Pancreatic Cancer: ABRAXANE 125 mg/m<sup>2</sup> on Days 1, 8 and 15 of each 28-day cycle. Recommended gemcitabine dose is 1000 mg/m<sup>2</sup> beginning immediately after the end of ABRAXANE administration on Days 1, 8 and 15 of each 28-day cycle.

Dose adjustments: Required for severe neutropenia, severe peripheral neuropathy and certain non-haematological toxicities (see full PI for details). Hepatic Impairment: Patients with severe hepatic impairment should not be treated with ABRAXANE. Consider dose reduction in patients with bilirubin >2 ULN. Refer to full PI for details.

**Administration:** Administered intravenously over 30 minutes. No premedication to prevent hypersensitivity reactions is required for ABRAXANE. Do not mix any other drugs with the ABRAXANE infusion.

Preparation for Intravenous Administration: Reconstitute with 20 mL of 0.9% Sodium Chloride. Inject appropriate amount of reconstituted ABRAXANE into an empty, sterile, polyvinyl chloride (PVC) or non-PVC type IV bag for IV infusion. Protect from light. For more details, refer to full PI.

Patients should consult their oncologist or the ABRAXANE Consumer Medicine Information available on [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

ABRAXANE<sup>®</sup> is a registered trademark of Celgene Corporation.

ABRAXANE<sup>®</sup> is distributed by STA under license from Celgene Corporation, in Australia and New Zealand.

## About Specialised Therapeutics Australia

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present

encompasses oncology, infectious diseases, and haematology. STA also has interests in the therapeutic areas of ophthalmology, respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

## References

1. ABRAXANE Product Information.
2. Von Hoff DD et al. Increased Survival in Pancreatic Cancer with nab-Paclitaxel plus Gemcitabine. *N Engl J Med* 2013; 369(18):1691-703.
3. Goldstein D et al. Oral Abstract # 178. Updated survival from a randomized phase III trial (MPACT) of nab-Paclitaxel plus gemcitabine versus gemcitabine alone for patients (pts) with metastatic adenocarcinoma of the pancreas. ASCO GI 2014.
4. Cancer in Australia. An Overview 2012. Australian Institute of Health and Welfare (AIHW).
5. Gradishar WJ et al. *J Clinical Oncology* 2005;23:7794-7803.

---

# **ARIAD Announces Commercial Agreement for ICLUSIG® (ponatinib) in Australia**

**Melbourne, Australia January 28, 2014:** ARIAD Pharmaceuticals, Inc. (NASDAQ: ARIA) and Specialised Therapeutics Australia Pty Ltd (STA), a specialty pharmaceutical company, today announced that ARIAD has granted STA exclusive rights to commercialize Iclusig® (ponatinib) in Australia in patients with Philadelphia-positive (Ph+) leukemias.

Under the terms of the agreement, STA will be responsible for obtaining marketing authorization and pricing and reimbursement approval of Iclusig and assisting ARIAD in regulatory filings for Iclusig in Australia. STA will book sales of Iclusig to pharmacies and other distributors, while ARIAD will supply packaged drug to STA. The term of the agreement is seven years from the first commercial sale of Iclusig following reimbursement approval. At the conclusion of the term, ARIAD will have the option to take over commercialization of Iclusig in Australia or to extend the agreement with STA.

“This agreement illustrates how we plan to make Iclusig available to patients in geographies where we do not anticipate setting up our own commercial activities near term,” said Marty J. Duvall, executive vice president and chief commercial officer of ARIAD. “STA has a proven track-record in oncology marketing and market access in Australia and is successfully distributing several important oncology brands in this region.”

ARIAD submitted a marketing application for Iclusig in the third quarter of 2013 to the Therapeutic Goods Administration (TGA) in Australia. Marketing approval and commercial launch of Iclusig are expected in the fourth quarter of 2014. Prior to launch, ARIAD and STA will collaborate to make Iclusig available to patients with refractory chronic myeloid leukemia (CML) and Ph+ acute lymphoblastic leukemia (ALL) under a Special Access Program.

“Iclusig is as an important cancer medicine for patients with difficult-to-treat CML or Ph+ ALL who have few options available to them,” said Carlo Montagner, chief executive officer at STA. “We look forward to a successful collaboration with ARIAD providing refractory CML patients in Australia with a new highly effective treatment option.”

According to the Australian Institute of Health and Welfare, there are more than 1,500 patients in Australia being treated for CML and approximately 290 patients are newly diagnosed with the disease each year.

“Some patients with this disease build resistance to current therapies and eventually run low on treatment options,” said Professor Timothy Hughes, Consulting Haematologist at the Royal Adelaide Hospital and one of the PACE trial investigators. “I anticipate that Iclusig will be a valuable new therapy for adult patients with refractory CML.”



## **About Iclusig® (ponatinib)**

Iclusig is a kinase inhibitor. The primary target for Iclusig is BCR-ABL, an abnormal tyrosine kinase that is expressed in chronic myeloid leukemia (CML) and Philadelphia-chromosome positive acute lymphoblastic leukemia (Ph+ ALL). Iclusig was designed using ARIAD's computational and structure-based drug design platform specifically to inhibit the activity of BCR-ABL. Iclusig targets not only native BCR-ABL but also its isoforms that carry mutations that confer resistance to treatment, including the T315I mutation, which has been associated with resistance to other approved TKIs.

## **About Specialised Therapeutics Australia**

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present encompasses oncology, hematology, ophthalmology and infectious diseases. STA also has interests in the therapeutic areas of respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

## **About ARIAD**

ARIAD Pharmaceuticals, Inc., headquartered in Cambridge, Massachusetts and Lausanne, Switzerland, is an integrated global oncology company focused on transforming the lives of cancer patients with breakthrough medicines. ARIAD is working on new medicines to advance the treatment of various forms of chronic and acute leukemia, lung cancer and other difficult-to-treat cancers. ARIAD utilizes computational and structural approaches to design small-molecule drugs that overcome resistance to existing cancer medicines. For additional

information, visit <http://www.ariad.com> or follow ARIAD on Twitter (@ARIADPharm).

This press release contains “forward-looking statements” which are based on management’s good-faith expectations and are subject to certain factors, risks and uncertainties that may cause actual results, outcome of events, timing and performance to differ materially from those expressed or implied by such statements. These factors, risks and uncertainties include, but are not limited to the Company’s ability to manufacture, and supply STA with Iclusig; the ability of STA to perform the contracted services, such as obtaining marketing authorization and pricing and reimbursement approval for Iclusig in Australia; STA’s ability to distribute, promote, market and sell Iclusig in Australia; the timing and scope of the marketing authorizations, as well as the level of pricing obtained in Australia; the availability of Iclusig to patients under a Special Access Program in Australia; third-party reimbursement; and the timing and success of sales of Iclusig in Australia. These factors, risks and uncertainties also include, but are not limited to: the costs associated with ARIAD’s development and manufacturing, commercial and other activities; the adequacy of capital resources and the availability of additional funding; and other factors detailed in the Company’s public filings with the U.S. Securities and Exchange Commission. The information contained in this press release is believed to be current as of the date of original issue. After the date of this document, the Company does not intend to update any of the forward-looking statements to conform to actual results or to changes in the Company’s expectations, except as required by law.

## **Contacts**

Carlo Montagner  
Chief Executive Officer  
Specialised Therapeutics Australia  
(03) 9859 1493

---

# ABC News 24: January 2014

The following story appeared on the ABC in July 2014 announcing a PBS listing.  
Click to view.



---

## Updated Pancreatic Cancer Data Presented at International Cancer Conference



**Melbourne, January 2014** - A leading breast cancer drug, ABRAXANE® (nanoparticle albumin-bound paclitaxel), has been shown to extend overall survival (OS) for patients with metastatic pancreatic cancer when used in combination with current standard of care, gemcitabine, with some patients surviving longer than three years.<sup>1</sup>

The updated OS data from the pivotal Phase III Metastatic Pancreatic Adenocarcinoma Clinical Trial (MPACT)<sup>1,2</sup> of ABRAXANE in combination with gemcitabine, was presented at the recent American Society of Clinical Oncology (ASCO) Gastrointestinal (GI) Conference in San Francisco by Australian Oncologist, Professor David Goldstein.

Professor Goldstein, who is based at the Prince of Wales Hospital in New South Wales, said that the data was “extremely encouraging” and paved the way for ABRAXANE to be used as a first-line therapy in the treatment of pancreatic cancer.

“Importantly, this large multinational trial in metastatic pancreatic cancer has demonstrated important three year survival rates, with 4% of patients in the

ABRAXANE plus gemcitabine arm alive after three years, compared to none in the gemcitabine alone arm, and a near 30% improvement in survival outcomes," he said.

"This cancer is the fourth most common cancer in Australia for both men and women<sup>3</sup> and typically patients diagnosed with metastatic pancreatic cancer have a median life expectancy of approximately three to six months. These patients have very limited treatment options as there have been no new medications approved for metastatic pancreatic cancer in nearly seven years.

Data from this important study shows that ABRAXANE when used in conjunction with standard of care gemcitabine can substantially improve OS, with predictable and manageable side effects."

The MPACT study was conducted at a number of sites internationally, including 20 sites in Australia, and involved 861 patients, of which 120 were Australian.

The median survival benefit in the updated analysis was extended in the ABRAXANE + gemcitabine arm with a 2.1 month OS improvement compared to gemcitabine alone (OS; median 8.7 months vs 6.6 months; HR=0.72;  $p<0.0001$ ).<sup>1</sup> This compares favourably with the 1.8 months improvement previously reported in the New England Journal of Medicine (OS; median 8.5 months vs 6.7 months; HR=0.72;  $p<0.00001$ ).<sup>2</sup>

The Chief Executive Officer of Australian biopharmaceutical company, Specialised Therapeutics Australia (STA), Mr Carlo Montagner, said ABRAXANE is currently approved in the United States and Europe as a first-line therapy in combination with gemcitabine for patients with metastatic pancreatic cancer.

He said a submission to the Therapeutic Goods Administration (TGA) for the regulatory approval of ABRAXANE (in combination with gemcitabine) for first-line treatment of locally advanced unresectable or metastatic adenocarcinoma of the pancreas was made in 2013. An application for a Pharmaceutical Benefits Scheme (PBS) listing for this indication has also been submitted with the Pharmaceutical Benefits Advisory Committee (PBAC) and will be reviewed at the March 2014 meeting.

Mr Montagner commented: "We are extremely pleased with these results

presented at the ASCO GI conference and look forward to a TGA approval for ABRAXANE and a subsequent PBS listing for Australian patients with this difficult to treat cancer.”

ABRAXANE is currently TGA approved for the treatment of metastatic breast cancer and first-line Non Small Cell Lung Cancer (NSCLC).<sup>4</sup>

## **About the MPACT Study<sup>2</sup>**

The MPACT study was a Celgene-sponsored, open-label, randomised, international study of 861 patients with metastatic pancreatic cancer. Patients were randomised to receive either ABRAXANE plus gemcitabine (125 mg/m<sup>2</sup> followed by 1000 mg/m<sup>2</sup> gemcitabine for 3 weeks followed by a week of rest) or gemcitabine alone (1000 mg/m<sup>2</sup> administered weekly for 7 weeks followed by a week of rest followed by cycles of weekly administration for 3 weeks followed by one week of rest).

The primary endpoint for the study was overall survival. Secondary endpoints were progression-free survival and overall response rate determined by independent radiological review. Other endpoints included progression-free survival and overall response rate determined by investigator, and the safety and tolerability of this combination in this patient population.

## **About Advanced Pancreatic Cancer**

Advanced pancreatic cancer is a difficult-to-treat cancer with the lowest survival rates among all cancer types. Across all patients with pancreatic cancer, relative 5-year survival is approximately 5% and is less than 2% for those with advanced disease.<sup>3</sup> There are two main types of pancreatic cancer - adenocarcinomas, which account for approximately 90% of all pancreatic cancer, and neuroendocrine tumours. Pancreatic cancer is relatively uncommon with new cases accounting for only 2.1% of all newly diagnosed cancers. However, pancreatic cancer is the fourth most common cause of cancer death for men and women in Australia, and the ninth most commonly diagnosed cancer in Australia.<sup>4</sup>

# About ABRAXANE®

ABRAXANE is a solvent-free, nanoparticle chemotherapy treatment option approved for the treatment of metastatic breast cancer and NSCLC.<sup>4</sup>

ABRAXANE is indicated for the treatment of metastatic carcinoma of the breast after failure of anthracycline therapy. ABRAXANE, in combination with carboplatin, is indicated for the first-line treatment of NSCLC in patients who are not candidates for potentially curative surgery and/or radiation.<sup>4</sup>

In Australia, ABRAXANE is currently listed on the PBS for the treatment of metastatic breast cancer and HER2 positive breast cancer in combination with trastuzumab. ABRAXANE is not listed on the PBS for the indication of NSCLC.

ABRAXANE is approved for metastatic breast cancer in over 40 countries including the U.S., Canada, European Union, Japan and China, and more than 180,000 cancer patients have received ABRAXANE therapy in the past five years.

ABRAXANE is currently in various stages of investigation for the treatment of the following cancers: metastatic melanoma, adjuvant pancreatic cancer, bladder cancer, ovarian cancer, and expanded applications for breast cancer.

Developed using the proprietary *nab*<sup>™</sup> technology platform, ABRAXANE is a nanoparticle protein-bound chemotherapy agent. ABRAXANE combines paclitaxel with albumin, a naturally-occurring human protein, to deliver the drug and eliminates the need for solvents in the administration process. Nanoparticle technology allows ABRAXANE to deliver a 49% higher dose compared to regular solvent-based paclitaxel without compromising safety and tolerability.<sup>45</sup>

## Contraindications and Side Effects<sup>4</sup>:

Like all medications, ABRAXANE may cause side effects.

ABRAXANE should not be used in patients who have baseline neutrophil counts of  $<1.5 \times 10^9$  /L.

Patients who have exhibited hypersensitivity reactions to ABRAXANE or human albumin should not be treated with ABRAXANE.

ABRAXANE is contraindicated during pregnancy and lactation.

Most common side effects ( $\geq 1/10$ ) caused by ABRAXANE include; neutropenia, anaemia, leucopenia, thrombocytopenia, lymphopenia, anorexia, peripheral neuropathy, hypoaesthesia, paraesthesia, nausea, diarrhoea, vomiting, constipation, stomatitis, alopecia, rash, arthralgia, myalgia, fatigue, asthenia, pyrexia. This is not a full list of all the side effects.

Before prescribing or for further information regarding ABRAXANE and potential side effects, physicians should review the ABRAXANE Product Information at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

Patients should consult their oncologist or the ABRAXANE Consumer Medicine Information available at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

ABRAXANE® is a registered trademark of Celgene Corporation.

ABRAXANE® is under license from Celgene Corporation and distributed by STA in Australia and New Zealand.

## **About Specialised Therapeutics Australia Pty Ltd**

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present encompasses oncology and infectious diseases. STA also has interests in the therapeutic areas of respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

- Study shows ABRAXANE plus gemcitabine combination therapy further extends overall survival benefit
- First Phase III trial in metastatic pancreatic cancer to report three year survival rates<sup>1</sup>
- New standard of care for patients with metastatic pancreatic cancer



## References:

1. Goldstein D et al. Oral Abstract # 178. Updated survival from a randomized phase III trial (MPACT) of nab-Paclitaxel plus gemcitabine versus gemcitabine alone for patients (pts) with metastatic adenocarcinoma of the pancreas. ASCO GI 2014.
  2. Von Hoff DD et al. Increased Survival in Pancreatic Cancer with nab-Paclitaxel plus Gemcitabine. N Engl J Med 2013; 369(18):1691-703.
  3. Cancer in Australia. An Overview 2012. Australian Institute of Health and Welfare (AIHW)
  4. ABRAXANE Product Information
  5. Gradishar WJ et al. J Clinical Oncology 2005;23:7794-7803
- 

# Specialised Therapeutics Australia Partners with Genomic Health Inc. to Deliver Novel Genomic Test to Breast Cancer Patients

**Melbourne, Australia January 7, 2014:** Australian women will potentially have greater access to the only genomic test validated to predict whether patients with early-stage invasive breast cancer would benefit from chemotherapy, following an agreement between Specialised Therapeutics Australia (STA) and Genomic Health, Inc. (NASDAQ: GHDX).

STA has struck an agreement to represent the important diagnostic technology known as the *Oncotype DX* Breast Cancer Assay from Genomic Health, the world's leading provider of genomic-based diagnostic tests that address both the overtreatment and optimal treatment of early stage cancer.

The *Oncotype DX* test is a 21 gene assay that predicts a patient's likely benefit from chemotherapy and the overall risk of breast cancer recurrence. This

technology has been shown to guide treatment decisions, sparing patients the impact of unnecessary chemotherapy while identifying those patients who may benefit from this additional treatment.

Under the terms of the agreement, STA will undertake all commercial operations including sales and marketing of the product within Australia as well as providing product support and practitioner education.

As part of this agreement, STA has partnered with Healthscope Pathology who will continue to oversee logistics in Australia, including tissue sample management.

Announcing the distribution agreement with Genomic Health, STA Chief Executive Officer Carlo Montagner said the *Oncotype DX* assay was a high calibre tool for women diagnosed with early stage breast cancer who sought to avoid chemotherapy where possible, because it provided information about the likelihood of a cancer recurrence.

“This ground breaking test, which has been universally adopted in the US, helps women make informed decisions,” he said.

“Many Australian women with early stage breast cancer have endured debilitating chemotherapy regimens as a precautionary measure. This test will arm women and their physicians with more information about the likelihood of the patient benefitting from chemotherapy, as well as recurrence, helping them make a well-informed treatment decision.”

Developed by US based Genomic Health, *Oncotype DX* has been evaluated in 15 clinical studies in more than 6,000 patients. These studies include a large validation study published in the *New England Journal of Medicine*<sup>1</sup> and a study published in the *Journal of Clinical Oncology* that examined whether *Oncotype DX* could predict the benefit of chemotherapy.<sup>2</sup> Since becoming available in 2004, more than 400,000 *Oncotype DX* tests have been requested by more than 19,000 physicians in over 70 countries.

The *Oncotype DX* Breast Cancer Test is appropriate for recently diagnosed women with Stage I or II node-negative, oestrogen-receptor-positive, HER2 negative, invasive breast cancer; and postmenopausal women with node-positive, hormone-receptor-positive, HER2 negative, invasive breast cancer. The

*Oncotype* DX results are provided in the form of a Recurrence Score<sup>®</sup>, a number between 0 and 100, which correlates to a specific likelihood of breast cancer recurrence within 10 years of initial diagnosis and the likely benefit of chemotherapy.

The *Oncotype* DX DCIS Breast Cancer Test is also appropriate for women with newly diagnosed pre-invasive or Ductal Carcinoma in Situ of the Breast (DCIS) who are treated with local excision, with or without adjuvant tamoxifen therapy. The DCIS Score<sup>™</sup> result predicts local recurrence of DCIS or invasive carcinoma and helps inform decisions regarding the need for additional treatments, like radiation, following surgical removal of the tumour.

The National Comprehensive Cancer Network (NCCN)<sup>3</sup>, the American Society of Clinical Oncology (ASCO)<sup>4</sup>, St Gallen<sup>5</sup>, and the European Society for Medical Oncology (ESMO)<sup>6</sup> have all incorporated the *Oncotype* DX test into their guidelines. In the UK, the National Institute for Health and Care Excellence (NICE) has recommended *Oncotype* DX as the only multi-gene breast cancer test for use in clinical practice to guide chemotherapy treatment decisions for patients with early-stage, hormone-receptor-positive, invasive breast cancer.<sup>7</sup>

“The *Oncotype* DX technology has played a critical role in predicting benefit of chemotherapy for more than 400,000 oestrogen-receptor-positive breast cancer patients globally in the past 10 years,” said Peter Zuendorf, Vice President, International, Genomic Health. “As adoption of *Oncotype* DX continues to grow, we are delighted to enter this partnership to broaden patient access to this unique test and the important information it provides to enable more individualised breast cancer treatment.”

This commercial arrangement between STA and Genomic Health commenced on 1st January 2014.

*Oncotype* DX, Recurrence Score and DCIS Score are trademarks of Genomic Health, Inc.

# About Specialised Therapeutics Australia

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present encompasses oncology and infectious diseases. STA also has interests in the therapeutic areas of respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

- Agreement to Distribute *Oncotype DX*® in Australia and New Zealand
- *Oncotype DX* now regarded as standard-of-care for early stage breast cancer treatment planning in the US since becoming available there in 2004

## References:

1. Paik et al. *NEJM* 2004; 351: 2817-2826
2. Paik et al. *J Clin Oncol* 2006; 24: 3726-3734
3. National Comprehensive Cancer Network® NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Breast Cancer. Version 3, 2013. Available at: [www.nccn.org](http://www.nccn.org)
4. Harris L et al. *J Clin Oncol* 2007; 25: 5287-5312
5. Goldhirsch A et al. *Ann Oncol* 2011; 22: 1736
6. Aebi S et al. *Ann Oncol* 2012; 22 (Supp 6): vi12-vi24
7. National Institute for Health and Care Excellence NICE diagnostic guidance 10. Issued September 2013. Available at: <http://guidance.nice.org.uk/DG10>

# Contacts

Carlo Montagner  
Chief Executive Officer  
Specialised Therapeutics Australia  
(03) 9859 1493

---

## **Specialised Therapeutics Australia Receives Therapeutic Goods Administration Approval for Brain Tumour Visualisation Drug - GLIOLAN®**

**Melbourne, Australia and Hamburg, Germany, November 2013:** A novel drug which assists neurosurgeons to better visualise and remove malignant brain tumours has been approved by the Therapeutic Goods Administration (TGA).

Until now, GLIOLAN (aminolevulinic acid HCl) has only been available via the Federal Government's Special Access Scheme (SAS). It will now be made widely available for use by neurosurgeons to treat patients with high grade glioma, specifically glioblastoma multiforme (GBM), which are tumours that typically have a very poor prognosis.

GLIOLAN is indicated in adult patients for **visualisation of malignant tissue during surgery for malignant gliomas that are glioblastoma multiforme (GBM) on preoperative imaging, and who are intended for resection of the**

## **tumour.**

GLIOLAN causes brain tumours (gliomas) to become fluorescent and glow during surgery. This enables neurosurgeons to better visualise these tumours and more completely remove them. GLIOLAN is given to the patient as a drink three hours before surgery. During surgery, a modified neurosurgical microscope fitted with a specialised blue operating light is used, which causes cancerous tissue to glow fluorescent red whilst normal brain tissue appears blue.

Melbourne bio-pharmaceutical company Specialised Therapeutics Australia Pty Ltd (STA) in-licenses the drug from German partner photonamic GmbH and Co. KG.

Announcing the TGA approval, STA chief executive officer Mr Carlo Montagner said GLIOLAN had already been used to treat over 100 Australian patients via the SAS and a number of hospitals have been quick to upgrade neurosurgical microscopes with fluorescence capability.

“We are pleased with the positive response from neurosurgeons since GLIOLAN was made available via the SAS and this approval from the TGA is an extremely positive outcome,” he said.

“It has always been our intention to make this high class compound available to all patients who may benefit. Brain tumour surgery using GLIOLAN has been widely adopted throughout Europe and we expect a similar uptake in Australia to improve outcomes for all GBM patients.”

The chief executive officer of photonamic Mr Ulrich Kosciessa said: “The approval in Australia is another milestone in our global development of GLIOLAN, which is now registered in more than 30 countries world wide.

“GLIOLAN was developed to provide neurosurgeons with an effective tool to increase radicality of brain tumour resection without compromising safety for the patients. We are pleased that our partner STA has successfully been able to achieve an approval from the TGA.”

International studies have shown that use of GLIOLAN during brain tumour surgery has nearly doubled the rate of achieving a complete resection, which in turn has resulted in a doubling of the number of patients without progression of

their brain cancer six months post surgery.<sup>1</sup>

The pivotal Phase III registration study published in The Lancet Oncology medical journal reported complete resection of the malignant brain tumour tissue was achieved in 65% of patients receiving GLIOLAN, compared to 36% of patients in the control arm. This resulted in 6-month progression-free survival being achieved in 41% of patients receiving GLIOLAN compared to 21.1% of patients who received surgery without the use of the drug.<sup>1</sup>

Brisbane neurosurgeon, Lindy Jeffree, has used GLIOLAN in 36 patients since the drug was first made available via the SAS. She regards fluorescence guided surgery as an important tool in helping surgeons distinguish parts of a tumour which would otherwise be invisible to the naked eye.

She commented: “It makes it much easier to distinguish tumour from normal brain tissue, which has undoubtedly assisted during some complex surgical procedures. Our aim is to provide optimal patient benefit. Using GLIOLAN to see tumour tissue more clearly enables better and more thorough resection which can make a big difference to a patient’s response to subsequent treatment and ultimately to survival.”

“I am extremely pleased to see this drug being made more widely available to improve surgical outcomes for patients with GBM around the country.”

The approval by the TGA approval brings the number of countries where GLIOLAN is registered to 31, including 27 in the EU as well as Japan, Korea and Taiwan. GLIOLAN was first approved in Europe in 2009 and is marketed by medac in Europe, Africa, South America and Asia (except Japan and Korea).

- Novel drug which improves visualisation and resection of malignant brain tumours now widely available
- Twice as many patients are without progression of brain cancer six months after surgery with GLIOLAN
- To date over 100 Australian patients have been treated with GLIOLAN via the Federal Government’s Special Access Scheme

The following Australian hospitals currently perform fluorescence-guided resection of brain tumours using GLIOLAN:

1. Royal Brisbane and Woman's Hospital, Queensland
2. The Wesley Hospital, Queensland
3. The Mater Private Hospital, Queensland
4. Princess Alexandra Hospital, Queensland
5. Prince of Wales Hospital, New South Wales
6. Newcastle Private Hospital, New South Wales
7. Calvary Hospital, Tasmania
8. The Royal Melbourne Hospital, Victoria
9. St Vincent's Private Hospital, Victoria

## **About GLIOLAN<sup>®</sup>**

The active substance in GLIOLAN, aminolevulinic acid (ALA), is a photoreceptive compound which is absorbed by cells in the body, where it is converted by enzymes into fluorescent chemicals, particularly protoporphyrin IX (PPIX). Since glioma cells take up more of the active substance and convert it more rapidly into PPIX, higher levels of PPIX accumulate in the cancer cells than in normal tissue. When illuminated under blue light of a specific wavelength, the PPIX in the tumour glows an intense red, while the normal brain tissue appears blue. This enables the surgeon to see the tumour more clearly during brain surgery and to remove it more accurately, sparing healthy brain tissue.<sup>2</sup>

Like all medications GLIOLAN may cause side effects. GLIOLAN should not be used in patients with hypersensitivity to ALA or porphyrins, or in cases of acute or chronic porphyria, or in pregnancy. Cardiac disorders, gastrointestinal disorders and skin and subcutaneous disorders are all reported as being uncommon.

## **About Specialised Therapeutics Australia**

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to



provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present encompasses oncology and infectious diseases. STA also has interests in the therapeutic areas of respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

## **About photonamic GmbH and Co KG**

photonamic GmbH and Co KG was established in 2003 to develop photosensitisers in the field of fluorescence guided diagnostics and photodynamic therapy. photonamic has developed ALA for the fluorescence guided resection of glioblastoma (GLIOLAN) and for the photodynamic therapy of skin lesions (ALACARE). Both products are approved in Europe and will further be developed for the global market. photonamic is based in Hamburg, Germany.

## **References:**

1. Stummer W, Pichlmeier U, Meinel T, et al., Fluorescence-guided surgery with 5-aminovulinec acid for resection of malignant glioma: a randomised controlled multicentre phase III trial, *Lancet Oncol*, 2006;7:392-401
2. European Public Assessment Report

## **Contacts**

Carlo Montagner  
Chief Executive Officer  
Specialised Therapeutics Australia  
(03) 9859 1493

---

# Melbourne Biopharmaceutical Company Named First Corporate Sponsor of International Medical Aid Organisation



**Melbourne, June 2013** Melbourne biopharmaceutical company Specialised Therapeutics Australia has become the first corporate sponsor of an international charitable organisation which enables medical and surgical specialists to impart their expertise to colleagues in developing countries.

The company has become the inaugural donor for the Specialists Without Borders (SWB) organisation, an Australian based non-profit entity established in 2005 to provide specialist medical and allied health teaching in developing countries.

The specialist teaching from some of Australasia's leading academic doctors, covers areas such as trauma surgery, orthopaedics, neurosurgery, anaesthetics, urology, radiology, oncology, paediatrics, perinatal health and psychiatry.

Announcing the \$50,000 donation, STA co-founders Carlo Montagner and Bozena Zembrzuski said this funding would immediately assist a specialised teaching seminar to be undertaken in Malawi and Zimbabwe in September 2013, with expansion of the teaching programme into Asia in 2014. STA will also provide communications and marketing support to SWB as part of its sponsorship.

20 Australian doctors and five select medical students will travel to each location for one week of teaching. Supported by four major Australian universities, the organisation has pioneered innovative teaching techniques in medicine and surgery and is rapidly becoming a valued medical teaching resource in developing countries. SWB has recently completed a five-year teaching module in Rwanda.

Ms Zembrzuski said these intensive teaching programs in developing countries were designed to empower local communities and reduce poverty by educating local doctors in cutting-edge medical developments and techniques which are often commonplace in the western world.

"STA is committed to improving health outcomes not only for Australians but for people in more remote parts of the world who may not have access to the same facilities and top-level expertise we are lucky enough to enjoy," she said.

"We hope this donation plays a part in helping SWB make real inroads in these developing countries, with the ultimate goal that doctors in these areas are equipped to give high class care to their own communities."

Working in full partnership with local communities and health care services, SWB co-ordinates an international body of medical specialists committed to providing medical education which enables sustainable health care solutions.

SWB chairman Dr Paul Anderson said all Australian doctors involved in the mission covered their own airfare and accommodation expenses.

"We are extremely pleased to accept this generous contribution from STA and are committed to ensuring these funds will help educate our colleagues in other parts of the world," Dr Anderson said.

“We can make a real difference. And it is this kind of support that will enable us to further develop and continue to provide programs addressing areas of high need in terms of medical education in the third world.

“Every trip we make brings us closer to achieving our goals of helping these communities help themselves with improved medical care and high class training.”

## **About Specialised Therapeutics Australia**

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present encompasses oncology and infectious diseases. STA also has interests in the therapeutic areas of respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)

## **About Specialists Without Borders**

Specialists Without Borders (SWB) is an Australia-based non-profit organisation contributing towards sustainable medical education in developing countries. SWB recognises the vital importance of information-sharing through the participation in global health, education and aid networks and works within the context of the United Nations Millennium Development Goals (MDGs) towards a more just, stable and secure world.

Through SWB, groups of international medical specialists volunteer their time to lead training seminars in developing countries. Each seminar is undertaken in full partnership with local communities, hospitals, health departments, medical schools and universities. Projects include, where possible, the upgrading of facilities and the provision of educational materials and textbooks to enhance medical learning and practice.

---

# New 'Superbug' Antibiotic Approved for Use in Australia

**MELBOURNE, Australia - April 26, 2013** - An effective new antibiotic designed to specifically treat the common superbug\* infection *Clostridium difficile*-associated diarrhoea will be available to patients in Australia from 14th May 2013.

Melbourne biopharmaceutical company Specialised Therapeutics Australia Pty Ltd (STA) has received Therapeutic Goods Administration (TGA) approval to market the drug DIFICID (fidaxomicin) in Australia. Until now, it has only been available in Australia under the Special Access Scheme.

DIFICID is indicated for the treatment of confirmed *Clostridium difficile* (CDI) infections in adults.<sup>1</sup>

The macrocyclic antibiotic therapy, taken in tablet form, is regarded as a breakthrough treatment to help fight serious CDI, which typically develops in patients following broad-spectrum antibiotic use. CDI targets the large intestine, causing diarrhoea which can range from moderate & debilitating to severe & life-threatening. It is extremely common in hospitals and aged care facilities as older patients are particularly vulnerable, and can be fatal.<sup>2</sup>

A recent media report indicated 14 Victorians died from the infection during a 15-month period in 2010 and 2011.<sup>3</sup> According to data generated by the Quebec provincial hospitalisation database, there were 7004 cases of *C. difficile* across Quebec from April 1st 2003 to March 31st 2004, and 1270 people died after contracting CDI.<sup>4</sup>

Medical experts say Australian infection rates have at least doubled in recent

years in major public hospitals, but concede the incidence of CDI is under reported.

STA Chief Executive Officer Mr Carlo Montagner is excited about the valuable treatment alternative DIFICID offers Australian patients who contract CDI.

“DIFICID is a potentially life saving drug for this extremely serious infection plaguing public hospitals and the wider community,” he said. “Unfortunately, it is estimated that almost 30% of patients can have a recurring infection. DIFICID is the only approved drug on the market which studies have shown will lower the risk of that infection returning.”

DIFICID is the first in a new class of antibiotics which are minimally absorbed by the bloodstream and have been shown to fight CDI while leaving healthy gut flora untouched.<sup>5</sup>

Hypervirulent strains of *C. difficile*, including the PCR ribotype 027 strain recently identified in Australia, have been associated with epidemic spread and high rates of severe disease and death.<sup>6</sup>

Risk factors for CDI include exposure to antimicrobial drugs, gastric acid-suppressive therapy, advanced age, prolonged hospitalisation, cancer chemotherapy, co-morbidity and immuno-suppression. Although most cases have been in hospital inpatients, increasing numbers of community-associated cases are now being reported.<sup>2</sup>

Leading Australian CDI expert Professor Thomas Riley from The University of Western Australia, acknowledged that studies had demonstrated patients treated with DIFICID were significantly less likely to develop recurrent infections.<sup>7,8</sup>

He regarded DIFICID as an important new treatment alternative, with infection rates of *C. difficile* climbing substantially in public hospitals around the country.

“Introducing DIFICID to Australia basically means we have another drug in the arsenal to treat this infection. Until now, we have had only two drugs available.

“Fewer recurrences will help contain the spread of the illness. Most importantly, DIFICID will benefit individual patients, who become weaker and more vulnerable

with each recurrent infection, enormously.”

STA licenses DIFICID for the Australian market from US-based Optimer Pharmaceuticals. Optimer Chief Executive Officer & Chairman of the Board, Dr Henry McKinnell, said he was confident DIFICID would provide a valuable new treatment option for an unmet medical need in Australia. “With the recent approval in Australia, fidaxomicin is now approved by four regulatory agencies, broadening access to patients in need across the globe,” said Dr. Henry McKinnell. “CDI infections represent a global healthcare challenge, and we believe an innovative drug like DIFICID that can deliver a substantial clinical improvement over existing therapies is an important new option that should be widely available to patients.”

## **About DIFICID<sup>®</sup>**

Fidaxomicin is a novel antibiotic agent and the first of a new class of antibacterials called macrocycles. Fidaxomicin is bactericidal against *C difficile* in vitro, inhibiting RNA synthesis by RNA polymerases.<sup>1</sup>

DIFICID was studied for the treatment of CDI in two randomised Phase III studies and was found to have equivalent efficacy to vancomycin. Notably, DIFICID was associated with significantly greater improvements in the rate of sustained clinical response and significantly lower rates of CDI recurrence (than vancomycin).<sup>1,7,8</sup>

## **Contraindications and side effects:<sup>1</sup>**

Like all medications, DIFICID may cause side effects. DIFICID should not be used in patients who are hypersensitive to any ingredient in the formulation or component of the container. As there is minimal systemic absorption of DIFICID, it should not be used for the treatment of systemic infections. Most common side effects ( $\geq 1/10$ ) caused by DIFICID include nausea, constipation and vomiting.

For further information regarding DIFICID and potential side effects, physicians

should review the DIFICID Approved Product Information available from [www.specialisedtherapeutics.com.au/index.php?q=clinician-resources.html](http://www.specialisedtherapeutics.com.au/index.php?q=clinician-resources.html) and patients should consult their prescribing physician or the DIFICID Consumer Medicine Information available in the pack or via [www.specialisedtherapeutics.com.au/index.php?q=dificid.html](http://www.specialisedtherapeutics.com.au/index.php?q=dificid.html)

## About CDI

CDI has become a significant medical problem in hospitals, long-term care facilities and the community. CDI is a serious illness resulting from infection of the inner lining of the colon by *C. difficile*, which produces toxins that cause inflammation of the colon, severe diarrhoea and, in the most serious cases, death. Patients typically develop CDI following the use of broad-spectrum antibiotics which disrupt normal gastrointestinal (gut) flora, possibly allowing *C. difficile* to enter the gut and flourish. Older patients in particular are at risk for CDI, potentially because of a weakened immune system or the presence of underlying disease. Approximately two-thirds of CDI patients are 65 years of age or older. Historically, approximately 20 - 30% of CDI patients who initially respond to treatment experience a clinical recurrence.<sup>7</sup>

## About Specialised Therapeutics Australia

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand. The STA therapeutic portfolio and pipeline at present encompasses oncology and infectious diseases. STA also has interests in the therapeutic areas of respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)



# About Optimer Pharmaceuticals

Optimer Pharmaceuticals, Inc. is a global biopharmaceutical company focused on developing and commercialising innovative hospital specialty products that have a positive impact on society. Optimer developed DIFICID (fidaxomicin) tablets, an FDA-approved macrolide antibacterial drug for the treatment of *Clostridium difficile*-associated diarrhoea (CDAD) in adults 18 years of age and older and is commercializing DIFICID in the US and Canada. Optimer also received marketing authorisation for fidaxomicin tablets in the European Union, where its partner, Astellas Pharma Europe, is commercialising fidaxomicin under the trade name DIFICLIR™. The company is exploring marketing authorisation in other parts of the world where *C. difficile* has emerged as a serious health problem. Additional information can be found at [www.optimerpharma.com](http://www.optimerpharma.com).

OPTIMER and DIFICID are trademarks of Optimer Pharmaceuticals, Inc. All other trademarks are the property of their respective owners.

\* Superbug is a common term to describe a bacterium that is resistant to multiple antibiotics.

## References:

1. DIFICID Approved Product Information
2. Cheng AC, Ferguson JK, Richards MJ, et al. Australasian Society for Infectious Diseases guidelines for the diagnosis and treatment of *Clostridium difficile* infection. *Med J Aust* 2011; 194: 353-358.
3. The Age, Saturday 26 May 2012.
4. Eggertson, L. *CMAJ* 2004; 171: (11) 1331-1332
5. Duggan ST. Fidaxomicin: In *Clostridium Difficile* Infection. *Drugs* 2011
6. Stuart R, Marshal C. *Clostridium difficile* infection: a new threat on our doorstep. *Med J Aust* 2011; 194: 331-332
7. Louie TJ, Miller MA, Mullane KM et al. Fidaxomicin versus Vancomycin for *Clostridium difficile* Infection *N Engl J Med* 2011;364:422-31.
8. Cornelly OA, Crook DW, Esposito R, et al. Fidaxomicin versus vancomycin for infection with *Clostridium difficile* in Europe, Canada, and the USA: a

double-blind, non-inferiority, randomised controlled trial. Lancet 2012

## Contacts

Carlo Montagner  
Chief Executive Officer  
Specialised Therapeutics Australia  
(03) 9859 1493

Emma Power  
Monsoon Communications  
(03) 9620 3333

---

# **ABRAXANE® Plus Gemcitabine Improves Survival in Phase III Study of Patients with Advanced Pancreatic Cancer**

**MELBOURNE, Australia - January 23, 2013** - Australian biopharmaceutical company Specialised Therapeutics Australia announces that a phase III clinical trial of world leading breast cancer drug ABRAXANE® (nanoparticle albumin-bound paclitaxel) in combination with current standard of care gemcitabine in patients with advanced pancreatic cancer has demonstrated substantially improved survival times, with double the number of patients surviving two years.<sup>1</sup>

The MPACT (Metastatic Pancreatic Adenocarcinoma Clinical Trial) investigation involved 861 treatment naïve patients internationally.

Researchers found those patients treated with ABRAXANE plus gemcitabine had a statistically significant improvement in overall survival compared to patients receiving gemcitabine alone .<sup>1</sup>

Moreover, ABRAXANE plus gemcitabine demonstrated a 59% increase in one-year survival (35% vs. 22%, p=0.0002) and demonstrated double the rate of survival at two years (9% vs. 4%, p=0.02) as compared to gemcitabine alone.<sup>1</sup>

ABRAXANE plus gemcitabine also demonstrated statistically significant improvements in key secondary endpoints compared to gemcitabine alone, including a 31% reduction in the risk of progression or death with a median progression-free survival (PFS) of 5.5 vs. 3.7 months (HR 0.69, P=0.000024) and an overall response rate (ORR) of 23% compared to 7% (response rate ratio of 3.19, p=1.1 x 10<sup>-10</sup>). Another endpoint assessed included time to treatment failure, which was significantly improved with the ABRAXANE combination compared to gemcitabine alone .<sup>1</sup>

“The past few decades have brought us very few treatment advances for patients with advanced pancreatic cancer, which is both deadly and incredibly difficult to treat with success,” said Daniel D. Von Hoff, M.D., F.A.C.P., Lead Principal Investigator of the MPACT study and Chief Scientific Officer for Scottsdale Healthcare’s Virginia G. Piper Cancer Centre Clinical Trials and Physician-In-Chief for TGen. “The fact that ABRAXANE plus gemcitabine demonstrated an overall survival benefit, and also did so at one and two years, is a significant step forward in offering potential new hope for our patients.”

Professor John Zalcborg, Chief Medical Officer and Executive Director of Cancer Medicine at the Peter MacCallum Cancer Centre in Melbourne, said the evidence strongly supported using ABRAXANE in combination with gemcitabine as a new standard of care to treat appropriate patients, many of whom were not diagnosed until the disease was metastatic.

While acknowledging that this advance could not be seen as a cure for pancreatic cancer, Professor Zalcborg said the 59% increase in the number of patients who lived beyond 12 months was very encouraging.

“We are extremely encouraged by the results of this study involving ABRAXANE

and regard this outcome as a significant breakthrough in terms of the future management of this disease," he said.

"In addition to treating women with metastatic breast cancer with ABRAXANE in the appropriate setting, we look forward to its approval in Australia for treating patients with advanced pancreatic cancer."

Specialised Therapeutics Australia (STA) Chief Executive Officer Mr Carlo Montagner said the positive data paved the way for Australian patients with advanced pancreatic cancer to access more effective treatment options.

He commented: "In Australia, pancreatic cancer is the fourth most common cause of death from cancer for both men and women<sup>2</sup> and very few treatment options exist for this group of patients. We are extremely pleased to demonstrate that ABRAXANE is capable of prolonging survival for patients with advanced pancreatic cancer and we hope to have ABRAXANE approved by the Australian Therapeutic Goods Administration (TGA) in the latter half of 2014."

The most common grade  $\geq 3$  treatment-related adverse events in the study for ABRAXANE plus gemcitabine vs. gemcitabine alone were neutropenia (38% vs. 27%), fatigue (17% vs. 7%), and neuropathy (17% vs. 1%). In the ABRAXANE plus gemcitabine arm, the median time to neuropathy improvement was 29 days. There was no difference in serious life threatening toxicity (4% in each arm).<sup>1</sup>

Further details of the study will be highlighted in a late-breaking oral presentation by Dr. Daniel D. Von Hoff:

Abstract: LBA #148: Final results of a randomized phase III study of weekly nab-paclitaxel plus gemcitabine versus gemcitabine alone in patients with metastatic adenocarcinoma of the pancreas. Friday, January 25th between 2:00 to 3:30 pm PST at the American Society of Clinical Oncology's (ASCO) 2013 Gastrointestinal Cancers Symposium in San Francisco, CA.

These results are from an investigational study. ABRAXANE is not approved for the treatment of advanced pancreatic cancer. Following TGA review and approval, STA will seek to have ABRAXANE included on the Pharmaceutical Benefits Scheme (PBS) for the reimbursement of ABRAXANE for advanced pancreatic cancer.

# About the MPACT Study<sup>1</sup>

In the MPACT (**M**etastatic **P**ancreatic **A**denocarcinoma **C**linical **T**rial) study, a Celgene-sponsored, open-label, randomised, international study of 861 patients with metastatic pancreatic cancer were randomised to receive either ABRAXANE plus gemcitabine (125 mg/m<sup>2</sup> followed by 1000 mg/m<sup>2</sup> gemcitabine for 3 weeks followed by a week of rest) or gemcitabine alone (1000 mg/m<sup>2</sup> administered weekly for 7 weeks followed by a week of rest followed by cycles of weekly administration for 3 weeks followed by one week of rest).

The primary endpoint for the study is improvement in overall survival. Secondary endpoints were progression-free survival, and overall response rate determined by independent radiological review. Other endpoints included progression-free survival, overall response rate determined by investigator and the safety and tolerability of this combination in this patient population.

## About Advanced Pancreatic Cancer

Advanced pancreatic cancer is a difficult-to-treat cancer with the lowest survival rates among all cancer types. Across all patients with pancreatic cancer, relative 5-year survival is 6% and is less than 2% for those with advanced disease. There are two main types of pancreatic cancer - adenocarcinomas, which accounts for approximately 90% of all pancreatic cancer, and neuroendocrine tumors. Pancreatic cancer is relatively uncommon with new cases accounting for only 2.1% of all newly diagnosed cancers. However, pancreatic cancer is the fourth most common cause of cancer death for men and women in the United States and Australia, and the ninth most commonly diagnosed cancer in Australia.<sup>2</sup>

## About ABRAXANE<sup>®</sup>

ABRAXANE is a solvent-free, nanoparticle chemotherapy treatment option for metastatic breast cancer.<sup>3</sup> In Australia, ABRAXANE is currently listed on the PBS

for the treatment of metastatic breast cancer and HER2 positive breast cancer in combination with trastuzumab.

ABRAXANE is approved for metastatic breast cancer in over 40 countries including the U.S., Canada, European Union, Japan and China, and more than 500,000 cancer patients have received ABRAXANE therapy in the past five years.

In Australia, ABRAXANE has been granted orphan drug designation by the Therapeutic Goods Administration for the treatment of pancreatic cancer. Orphan drug status is granted to drugs used to treat relatively rare diseases such as pancreatic cancer and may allow for priority evaluation by the TGA.

ABRAXANE is currently in various stages of investigation for the treatment of the following cancers: metastatic melanoma, bladder, ovarian, and expanded applications for breast cancer.

Developed using the proprietary *nab*<sup>™</sup> technology platform, ABRAXANE is a nanoparticle protein-bound chemotherapy agent. ABRAXANE combines paclitaxel with albumin, a naturally-occurring human protein, to deliver the drug and eliminates the need for solvents in the administration process. Nanoparticle technology allows ABRAXANE to deliver a 49% higher dose compared to regular solvent-based paclitaxel without compromising safety and tolerability.<sup>3-4</sup>

In a randomised phase III study of metastatic breast cancer patients, ABRAXANE demonstrated nearly double the overall tumour response rate compared to solvent-based paclitaxel.<sup>3-4</sup>

Anthracycline pre-treated patients in the study lived significantly longer.<sup>5</sup> The tolerability with ABRAXANE and solvent-based paclitaxel was comparable, despite the 49% greater dose of paclitaxel administered as ABRAXANE.<sup>3-4</sup> Neutropenia was lower with ABRAXANE compared to solvent-based paclitaxel, although there was an increase in incidence of grade 3 peripheral neuropathy with ABRAXANE. However the median time to improvement, from grade 3 peripheral neuropathy to grade 2 or lower, was 22 days. No adverse events were reported that were not already known for paclitaxel.<sup>3-4</sup>

## Contraindications and side effects<sup>3</sup>:

Like all medications, ABRAXANE may cause side effects.

ABRAXANE should not be used in patients who have baseline neutrophil counts of  $<1.5 \times 10^9$  /L.

In patients who have exhibited hypersensitivity reactions to paclitaxel or albumin, patients should not be treated with ABRAXANE.

ABRAXANE is contraindicated during pregnancy and lactation.

Most common side effects ( $\geq 1/10$ ) caused by ABRAXANE include; neutropenia, anemia, leucopenia, thrombocytopenia, lymphopenia, anorexia, peripheral neuropathy, hypoaesthesia, paraesthesia, nausea, diarrhoea, vomiting, constipation, stomatitis, alopecia, rash, arthralgia, myalgia, fatigue, asthenia, pyrexia.

For further information regarding ABRAXANE and potential side effects, physicians should review the ABRAXANE Product Information and patients should consult their oncologist or the ABRAXANE Consumer Medicine Information available on [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au).

ABRAXANE<sup>®</sup> is a registered trademark of Celgene Corporation.

ABRAXANE<sup>®</sup> is distributed by STA under license from Celgene Corporation in Australia and New Zealand.

About Specialised Therapeutics Australia, Pty Ltd

Specialised Therapeutics Australia Pty Ltd (STA) is a biopharmaceutical company dedicated to working with leading pharmaceutical companies worldwide to provide acute care therapies for high unmet medical needs to people living in Australia and New Zealand.

Currently STA markets two world leading cancer and cancer supportive care therapies, ABRAXANE<sup>®</sup> (nab-paclitaxel) and ALOXI<sup>®</sup> (palonosetron HCl) respectively, and has recently licensed two new agents from the Helsinn Group. Firstly Anamorelin, which is a novel ghrelin receptor agonist for the treatment of anorexia-cachexia in NSCLC, and a fixed-dose combination product (in both oral and intravenous forms) containing netupitant, a neurokinin-1 (NK1) receptor antagonist, combined with Aloxi, a serotonin-3 (5-HT<sub>3</sub>) receptor antagonist. STA

also has interests in the therapeutic areas of anti-infectives with the rights to commercialise DIFICID® (fidaxomicin) for the treatment of Clostridium difficile infections, respiratory, dermatology, endocrinology and central nervous system (CNS). Additional information can be found at [www.specialisedtherapeutics.com.au](http://www.specialisedtherapeutics.com.au)


- ABRAXANE plus gemcitabine demonstrated highly statistically significant and clinically meaningful results across primary and key secondary endpoints and patient subgroups
- ABRAXANE plus gemcitabine patients showed 59% higher chance of survival at one year; survival rates doubled at two years
- A new standard of care for patients with advanced pancreatic cancer
- Oral Presentation Scheduled for Friday, January 25th at ASCO's Gastrointestinal Cancers Symposium Annual Meeting

### **References:**

1. Von Hoff DD et al. Abstract: LBA #148: Final results of a randomized phase III study of weekly nab-paclitaxel plus gemcitabine versus gemcitabine alone in patients with metastatic adenocarcinoma of the pancreas. ASCO GI 2013
  2. Cancer in Australia. An Overview 2012. Australian Institute of Health and Welfare (AIHW)
  3. Abraxane Product Information
  4. Gradishar WJ et al. J Clinical Oncology 2005;23:7794-7803
  5. Vukelja SJ et al. ASCO 2008, Abstract 1082
-



# ABRAXANE Meets Primary Endpoint of Overall Survival Improvement in Phase III Advanced Pancreatic Cancer Study



**Melbourne, November 2012** Specialised Therapeutics Australia (STA) is pleased to announce that the phase III, randomised, international study of ABRAXANE® (nanoparticle albumin-bound paclitaxel) in combination with gemcitabine in treatment-naïve patients with advanced pancreatic cancer met its primary endpoint of overall survival.

This is now the fourth disease in which ABRAXANE has demonstrated positive phase III results, including difficult to treat diseases where other taxanes have shown limited activity.

In the MPACT (Metastatic Pancreatic Adenocarcinoma Clinical Trial) study, a Celgene-sponsored, open-label, randomised, international study, 861 metastatic pancreatic cancer patients were randomised to receive either ABRAXANE plus gemcitabine (125 mg/m<sup>2</sup> followed by 1000 mg/m<sup>2</sup> gemcitabine for 3 weeks followed by a week of rest) or gemcitabine alone (1000 mg/m<sup>2</sup> administered weekly for 7 weeks followed by a week of rest followed by cycles of weekly administration for 3 weeks followed by one week of rest).

The primary endpoint for the study was improvement in overall survival. Secondary endpoints included evaluation of progression-free survival, objective tumour response and the safety and tolerability of the ABRAXANE/gemcitabine combination in this patient population.

The safety profile of ABRAXANE in combination with gemcitabine observed in the study was comparable with other ABRAXANE clinical trials in pancreatic cancer.

A late-breaker abstract has been submitted to the American Society of Clinical Oncology's (ASCO) 2013 Gastrointestinal Cancers Symposium being held in San Francisco on January 24-26, 2013, after which more details will be made available.

Australia contributed the second highest number of patients to this study with 120 patients enrolled. Combined with over 450 patients with advanced pancreatic cancer who have accessed ABRAXANE via STA's compassionate access program, substantial local clinical experience with ABRAXANE has been generated in this disease.

Based on the results of the MPACT study, STA look forward to submitting an application in 2013 to the Therapeutic Goods Administration and Pharmaceutical Benefits Advisory Committee for regulatory and PBS approval.

Please note these results are from an investigational phase III study. ABRAXANE is not currently approved for the treatment of advanced pancreatic cancer in Australia.

Before prescribing ABRAXANE please refer to the [Abraxane Product Information](#).  
For further information please contact STA on 1300 798 820.